

IHCAb™ CD45 (LCA) (BGT-CD45) mouse mAb

Cat #: B-IMW6156

Size: 100 uL

Storage: Store at -20°C. Avoid repeated freeze / thaw cycles.

Background

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be

signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitosis, and

oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two

tandem intracytoplasmic catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be

an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with

components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen

receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling.

Alternatively spliced transcripts variants of this gene

Product Information

Applications/Dilution: IHC-p 1:100-500, WB 1:200-1000;IF(paraffin section);Flow Cyt 1:50-200

Isotype/Source: Mouse, Monoclonal/IgG2b, Kappa

Specificity: This antibody detects endogenous levels of human CD45 (LCA). Heat-induced epitope retrieval (HIER)

Citrate buffer of pH6.0 was highly recommended as antigen repair method in paraffin section

Subcellular Location: Cell membrane; Single-pass type I membrane protein. Membrane raft. Colocalized with DPP4 in

membrane rafts

Expression: Isoform 1: Detected in thymocytes. Isoform 2: Detected in thymocytes. Isoform 3: Detected in thymocytes.

Isoform 4: Not detected in thymocytes. Isoform 5: Detected in thymocytes. Isoform 6: Not detected in thymocytes.

Isoform 7: Detected in thymocytes. Isoform 8: Not detected in thymocytes.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide

Biogradetech



Storage: Store at -15°C to -25°C




Fig.1. Human diffuse large B-cell lymphoma tissue was stained with Anti-CD45 Antibody.




Fig. 2. Human diffuse large B-cell lymphoma tissue was stained with Anti-CD45 Antibody.

Note:

The product listed herein is for research use only and is not intended for use in human or clinical diagnosis.

