

EZMeta™ Aconitase (ACO) Colorimetric Activity Kit

Cat #: D-AKC1290

Size: 48T / 96T

Storage: Stored at -20°C for 6 months, protected from light

Product Information

Applicable samples: Animal and Plant Tissues, Cells, Serum, Plasma

Assay Principle

Aconitase (ACO), an enzyme in the tricarboxylic acid cycle, catalyzes the conversion of citric acid to isocitrate. Citric acid itself is not easy to be oxidized. Under the action of aconitase, the hydroxyl group is transferred from the β carbon atom to the α carbon atom through the reaction of dehydration and water addition to generate isocitric acid that is easy to deoxidize, for further oxidative decarboxylation reaction be prepared. EZMeta™ Aconitase (ACO) Colorimetric Activity Kit provides a simple, convenient and rapid ACO activity detection method, which is suitable for the detection of animal tissues, plant tissues, cells, serum, plasma and other samples. The principle is that ACO catalyzes the conversion of citric acid into isocitrate, and the oxidative decarboxylation of isocitrate reduces NAD^+ to NADH, resulting in an increase in light absorption at 340 nm

Materials Supplied and Storage Conditions

Kit components	Size		Storage conditions
	48 T	96 T	
Extraction Buffer	50 mL	100 mL	4°C
Reagent I	10 mL	20 mL	4°C
Reagent II	0.75 mL	1.5 mL	-20°C, protected from light

Reagent III	10 mL	20 mL	4°C
Reagent IV	1	1	-20°C, protected from light
Reagent V	1	1	-20°C, protected from light
Reagen VI	1	1	-20°C, protected from light

Materials Required but Not Supplied

- Microplate reader or ultraviolet spectrophotometer capable of measuring absorbance at 340 nm
- 96-well UV plate or microquartz cuvette, precision pipettes, disposable pipette tips
- Ice maker, refrigerated centrifuge, incubator
- Deionized water
- Homogenizer (for tissue samples)

Reagent Preparation

Extraction Buffer: Ready to use as supplied. Equilibrate to room temperature before use. Store at 4°C.

Reagent I : Ready to use as supplied. Equilibrate to room temperature before use. Store at 4°C.

Reagent II : Ready to use as supplied. Equilibrate to room temperature before use. Store at -20°C, protected from light.

Reagent III: Ready to use as supplied. Equilibrate to room temperature before use. Store at 4°C.

Reagent IV: Prepare before use. For 48 T, add 2.5 mL deionized water, and for 96 T, add 5 mL deionized water, mix well.

The remaining reagents should be stored at -20°C and protected from light after aliquoting to avoid repeated freezing and thawing.

Reagent V: Prepare before use. For 48 T, add 0.75 mL of deionized water, and for 96 T, add 1.5 mL of deionized water, mix well. The remaining reagents should be stored at -20°C and protected from light after aliquoting to avoid repeated freezing and thawing.

Working Reagent VI: Prepare before use. For 48 T, add 6 mL Reagent III to Reagent VI, and for 96 T, add 12 mL Reagent III to Reagent VI, mix well. The remaining reagents should be stored at -20°C and protected from light after aliquoting to avoid repeated freezing and thawing.

Working Solution: The solution was freshly prepared just before use. For 48 T, add 0.5 mL deionized water, 0.5 mL Reagent III, 0.5 mL Reagent IV, and 0.5 mL Reagent V to 6 mL Working Reagent VI; for 96 T, add 1 mL deionized water, 1 mL Reagent III, 1 mL Reagent IV, and 1 mL Reagent V to 12 mL Working Reagent VI, and mix well for use. The remaining reagents should be stored at -20°C and protected from light after aliquoting to avoid repeated freezing and thawing for one week.

Sample Preparation

Note: Fresh samples are recommended, If not assayed immediately, samples can be stored at -80°C for one month. All samples and reagents should be on ice to avoid denaturation and deactivation.

1. Plasma and Serum: Direct detection.

2. Extraction of cytoplasmic protein and mitochondrial protein from cells and tissue:

(1) Weigh 0.1 g tissue or collect 5×10^6 cells, add 1 mL Extraction Buffer and 10 μ L Reagent II, homogenize on ice. Centrifuge at 600 g for 5 min at 4°C. Collect the supernatant to a new centrifuge tube and discard the pellet.

(2) Centrifuge the supernatant again at 11,000 g for 10 min at 4°C, and obtain the supernatant and precipitate respectively.

(3) (Optional) The supernatant collected in step 2 is cytoplasmic extract, which can be used to directly determine ACO leaking from mitochondria.

(4) Add 200 μ L Reagent I and 2 μ L Reagent II to the precipitate collected in step 2, resuspend the precipitate sufficiently, and use it to detect the activity of ACO in the next step.

Note: In order to guarantee the accuracy of experimental results, need to do a pre-experiment with 2-3 samples. If the samples are placed for a long time, different operation habits or other reasons, which resulted in a large amount of mMDH in the supernatant, then the supernatant must be tested.

Assay Procedure

1. Preheat the microplate reader or ultraviolet spectrophotometer for more than 30 min, and adjust the wavelength to 340 nm, ultraviolet spectrophotometer was returned to zero with deionized water.

2. Incubate Working Solution for 10 min at 37°C (mammal) or 25°C (other species).
3. Add 40 µL of sample, 160 µL of Working Solution in a 96-well UV plate or microquartz cuvette. After mixing quickly, record the absorbance values of 20 s and 3 min 20 s at 340 nm with a microplate reader, mark as A₁ and A₂, and calculate $\Delta A = A_2 - A_1$.

Note: If the sample absorbance value ΔA is greater than 0.5, the sample can be appropriately diluted with Reagent I, the calculated result multiplied by the dilution factor. It is not suggested to test too many samples at the same time, because enzyme activity is calculated by the variation of absorbance value per unit time.

Data Analysis

Note: We provide you with calculation formulae, including the derivation process and final formula. The two are exactly equal. It is suggested that the concise calculation formula in bold is final formula.

A. 96-well UV plates calculation formula

1. Calculation of ACO activity in serum (plasma)

Unit definition: one enzyme activity unit defines as 1 nmol NADH produced by each mL of serum (plasma) per min in the reaction system.

$$\text{ACO activity (U/mL)} = [\Delta A \times V_{\text{total}} \div (\epsilon \times d) \times 10^9] \div V_{\text{sample}} \div T = \mathbf{535.90 \times \Delta A}$$

2. Calculated by fresh weight of samples

Unit definition: one enzyme activity unit defines as 1 nmol NADH produced by 1 g tissue per min in the reaction system.

$$\text{ACO}_{\text{Supernatant}} \text{ activity (U/g fresh weight)} = [\Delta A_{\text{Supernatant}} \times V_{\text{Total}} \div (\epsilon \times d) \times 10^9] \div (V_{\text{Sample}} \div V_{\text{Extraction}} \times W) \div T = \mathbf{541.26 \times \Delta A_{\text{Supernatant}} \div W}$$

$$\text{ACO}_{\text{Pellet}} \text{ activity (U/g fresh weight)} = [\Delta A_{\text{Pellet}} \times V_{\text{Total}} \div (\epsilon \times d) \times 10^9] \div (V_{\text{Sample}} \div V_{\text{Total Sample}} \times W) \div T = \mathbf{108.25 \times \Delta A_{\text{Pellet}} \div W}$$

$$\text{Total ACO activity (U/g fresh weight)} = \text{ACO}_{\text{Supernatant}} \text{ activity} + \text{ACO}_{\text{Pellet}} \text{ activity} = \mathbf{541.26 \times \Delta A_{\text{Supernatant}} \div W + 108.25 \times \Delta A_{\text{Pellet}} \div W}$$

3. Calculated by cell density

Unit definition: one enzyme activity unit defines as 1 nmol NADH produced by 10⁴ cells per minute in the reaction system.

$$\text{ACO activity (U/10}^4 \text{ cells)} = [\Delta A \times V_{\text{Total}} \div (\epsilon \times d) \times 10^9] \div (V_{\text{Sample}} \div V_{\text{Total Sample}} \times 500) \div T = \mathbf{0.216 \times \Delta A}$$

Where: V_{Total}: total reaction volume, 2×10⁻⁴ L; ϵ : NADH molar extinction coefficient, 6.22×10³ mol/L/cm; d: 0.5 cm;

V_{Sample} : sample volume added, 0.04 mL; T: reaction time, 3 min; $\Delta A_{\text{Supernatant}}$: OD value of supernatant; $V_{\text{Extraction}}$: sample extract volume, 1.01 mL; W: sample weight, g; ΔA_{Pellet} : OD value of pellet; $V_{\text{Total Sample}}$: the volume of adding Reagents I and II, 0.202 mL; 500: total number of cells, 5×10^6 .

B. Microquartz cuvette calculation formula

The optical diameter d: 0.5 cm in the above calculation formula can be adjusted to d: 1 cm for calculation.

Disclaimer

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.