

## Recombinant Swine, IL10 / Interleukin-10,

### Animal Free & Carrier Free

Cat #: C-CYP120

Size: 5µg, 20µg

Shipping: Blue Ice

### Product Overview

Interleukin 10 (IL-10), also known as human cytokine synthesis inhibitory factor (CSIF), is an antiinflammatory cytokine. In humans, interleukin 10 is encoded by the IL10 gene. IL-10 signals through a receptor complex consisting of two IL-10 receptor-1 and two IL-10 receptor-2 proteins. Consequently, the functional receptor consists of four IL-10 receptor molecules. IL-10 binding induces STAT3 signalling via the phosphorylation of the cytoplasmic tails of IL-10 receptor 1 + IL-10 receptor 2 by JAK1 and Tyk2 respectively.

### Product Information

**Source:** *Escherichia Coli*.

**Purity:** >98% as determined by SDS-PAGE. Ni-NTA chromatography.


**Biological Activity:** Measure by its ability to induce proliferation in MC/9 -2 cells. The ED50 for this effect is < 5 ng/mL

**Endotoxin:** < 0.01 EU per 1 µg of the protein by the LAL method.

**Amino acid sequence:**

MSIKSENSCIHFPTSLPHMLRELRAAFGPVKSFFQTKDQMGDLLLTGSLLEDFKGYLGCQALSEMIQFYLEDVMPKAESDGEDIKEHVNSLG  
EKLKTLRLRLRRCHQFLPCENKSKAVEEVKSAFSKLQERGVYKAMGEFDIFINYIEAYMTMKMRKN with polyhistidine tag at the C-terminus

**Formulation:** Lyophilized from a sterile filtered aqueous solution in 1xPBS, pH 8.0.



SDS-PAGE analysis of recombinant swine IL-10

## Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H<sub>2</sub>O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

## Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

## Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.