

Recombinant Swine, IL15 / Interleukin-15,

Animal Free & Carrier Free

Cat #: C-CYP123 Size: 5µg, 20µg Shipping: Blue Ice

Product Overview

Interleukin-15 (IL-15) is a cytokine with structural similarity to Interleukin-2 (IL-2). Like IL-2, IL-15 binds to and signals through a complex composed of IL-2/IL-15 receptor beta chain (CD122) and the common gamma chain (gamma-C, CD132). IL-15 is secreted by mononuclear phagocytes (and some other cells) following infection by virus(es). This cytokine induces cell proliferation of natural killer cells; cells of the innate immune system whose principal role is to kill virally infected cells.

Product Information

Source: Escherichia Coli.

Purity: >95% as determined by SDS-PAGE. Ni-NTA chromatography.

Biological Activity: Measure by its ability to induce proliferation in TF-1 cells. The ED₅₀ for this effect is < 5.5 ng/mL

Endotoxin: < 0.01 EU per 1 μ g of the protein by the LAL method.


Amino acid sequence:

TWQHVISDLKKIEDLIRSIHMDATLYTESDAHPNCKVTAMKCFLLELRVILQESRNSDISDTVENLIILANSSLSSIEYKTESGCKECEELEEKNIN EFLKSFIHIVQMFINPS with polyhistidine tag at the C- terminus

Formulation: Lyophilized from a sterile filtered aqueous solution in 1×PBS, pH 7.4.







SDS-PAGE analysis of recombinant swine IL-15

Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.

2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H_2O . Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.

3. The reconstituted solution can be stored at 2-8°C for up to 1 week.

4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 μ g/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.

