

Recombinant Swine, IFN γ / Interferon gamma,

Animal Free & Carrier Free

Cat #: C-CYP135

Size: 5 μ g, 20 μ g

Shipping: Blue Ice

Product Overview

Interferon- γ is an effective multifunctional cytokine which is released mainly by activated NK cells and T cells. IFN- γ is primarily characterized based on its anti-viral activities, and then been proved to have several functions such as anti-proliferative, immune regulatory, and pro-inflammatory activities. IFN- γ can upregulate expression of MHC class I and II antigen by antigen-presenting cells.

Product Information

Source: *Escherichia Coli*.

Purity: >95% as determined by SDS-PAGE. Ni-NTA chromatography.

Biological Activity: Measure by its ability to protect PK15 cells infected with encephalomyocarditis(EMC) virus.


The ED₅₀ for this effect is < 40 pg/ mL.

Endotoxin: <0.01 EU per 1 μ g of the protein by the LAL method.

Amino acid sequence:

MQAPFFKEITILKDYFNASTSDVPNGGPLFLEILKNWKEESDKKIIQSQIVSFYFKFFEIFKDNQAIQRSMDVIKQDMFQRFLNGSSGKLNDF
EKLIKIPVDNLQIQRKAISELIKVMNDLSPRSNLRKRKRSQTMFQGQRASK with polyhistidine tag at the C-terminus

Formulation: Lyophilized from a sterile filtered aqueous solution in 1 \times PBS, pH 7.4.



SDS-PAGE analysis of recombinant swine IFN gamma

Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H₂O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.