

## Recombinant Human, IL11 / Interleukin-11,

### Animal Free & Carrier Free

Cat #: C-CYP136

Size: 5µg, 20µg, 100µg, 500µg, 1mg

Shipping: Blue Ice

### Product Overview

IL-11 is a cytokine and first isolated in 1990 from bone marrow-derived fibrocyte-like stromal cells. It was initially thought to be important for hematopoiesis, notably for megakaryocyte maturation, but subsequently shown to be redundant for platelets, and for other blood cell types, in both mice and humans. It is also known under the names adipogenesis inhibitory factor (AGIF) and was developed as a recombinant protein (rhIL-11) as the drug substance oprelvekin

### Product Information

**Source:** *Escherichia Coli*.

**Purity:** >98% as determined by SDS-PAGE. Ni-NTA chromatography.

**Biological Activity:** Measure by its ability to induce T11 cells proliferation. The ED<sub>50</sub> for this effect is < 0.2 ng/mL. The specific activity of recombinant human IL-11 is approximately >1 x 10<sup>7</sup> IU/mg.

**Endotoxin:** <0.1 EU per 1 µg of the protein by the LAL method.


**Amino acid sequence:**

PGPPPGPPRVSPDPRAELDSTVLLTRSLADTRQLAAQLRDKFPADGDHNLDSLPTLAMSAGALGALQLPGVLTRLRA

DLLSYLRHVQWLRRRAGGSSKLTLEPELGTQLARLDRLRLQLLMSRLALPQPPDPAPPPLAPPSSAWGGIRAAHAILGGLHLTDWAVR

GLLLLKTRL with polyhistidine tag at the N-terminus

**Formulation:** Lyophilized from a sterile filtered aqueous solution in 1×PBS, pH 7.4.



SDS-PAGE analysis of recombinant human IL-11

## Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H<sub>2</sub>O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

## Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

## Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.