

Recombinant Swine, FLT3L / Fms-Related Tyrosine Kinase 3 Ligand, Animal Free & Carrier Free

Cat #: C-CYP138

Size: 5µg, 20µg

Shipping: Blue Ice

Product Overview

Fms-related tyrosine kinase 3 ligand (FLT3LG) is a protein which in humans is encoded by the FLT3LG gene. FLT3 ligand is a receptor for the fl cytokine has a tyrosine-protein kinase activity & a growth factor that regulates proliferation of early hematopoietic cells. Flt3-Ligand synergizes with other CSFs and interleukins to induce growth and Differentiation

Product Information

Source: *Escherichia Coli*.

Purity: >95% as determined by SDS-PAGE. Ni-NTA chromatography.

Biological Activity: Measure by its ability to induce proliferation in OCI-AML5 cells. The ED₅₀ for this effect is < 5 ng/mL

Endotoxin: <0.1 EU per 1 µg of the protein by the LAL method.


Amino acid sequence:

MSPDCSFPHSPISSTFANTIRQLSDYLLQDYPVTVASNLQDDELCGAFWRLVLAQRWVGQLKTVAGSQMQKLEAVNTEIVFVTSCALQP

LPSCLRFVQANISHLLQDTSQQQLVALKPWITRRNFSRCLELQCQDPSTLLPPRSPGALEATSLPAPQASLLLLLLLLLPAALLL with

polyhistidine tag at the C-terminus

Formulation: Lyophilized from a sterile filtered aqueous solution in 1×PBS, pH 7.4.



SDS-PAGE analysis of of recombinant swine Flt-3 Ligand

Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H₂O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.