

## Recombinant Human, IL17F / Interleukin-17F,

### Animal Free & Carrier Free

Cat #: C-CYP158

Size: 5µg, 20µg, 100µg, 500µg, 1mg

Shipping: Blue Ice

### Product Overview

The protein encoded by this gene is a cytokine that shares sequence similarity with IL17. This cytokine is expressed by activated T cells, and has been shown to stimulate the production of several other cytokines, including IL6, IL8, and CSF2/GM-CSF. This cytokine is also found to inhibit the angiogenesis of endothelial cells and induce endothelial cells to produce IL2, TGFβ1/TGFβ, and monocyte chemoattractant protein-1.

### Product Information

**Source:** *Escherichia Coli*.

**Purity:** >98% as determined by SDS-PAGE. Ni-NTA chromatography.


**Biological Activity:** Measure by its ability to induce IL-6 secretion in 3T3 cells. The ED<sub>50</sub> for this effect is < 20 ng/mL.

**Endotoxin:** <0.1 EU per 1 µg of the protein by the LAL method.

**Amino acid sequence:**

MRKIPKVGHTFFQKPESCPPVPGGSMKLDIGIINENQRVSMRNIERSRSTSPWNYTVTWDPNRYPSEVVQAQCRNLGCINAQGKEDISM  
NSVPIQQETLVVRRKHQGCVSFQLEKVLVTVGCTCVTPVIHHVQ with polyhistidine tag at the C-terminus

**Formulation:** Lyophilized from a sterile filtered aqueous solution in 20 mM sodium acetate, pH 4.0.



SDS-PAGE analysis of recombinant human IL-17F

## Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H<sub>2</sub>O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

## Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

## Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.