

Recombinant Swine, CXCL13 / C-X-C Motif Chemokine Ligand 13, Animal Free & Carrier Free

Cat #: C-CYP175

Size: 5µg, 20µg

Shipping: Blue Ice

Product Overview

CXCL13, also known as BCA-1 (B Cell- Attracting chemokine 1) or BLC, is a recently identified new CXC chemokines.

This chemokine is expressed in the stomach, spleen, liver, appendix and lymph nodes. CXCL13 can only elicit its activities through CXCR5 receptor. BCA-1/BLC is a potent chemoattractant for B lymphocytes, and induces weak chemotactic response in T cells and macrophages. It should be noticed that CXCL13 shows no activity on neutrophils and monocytes.

Product Information

Source: *Escherichia Coli*.

Purity: >98% as determined by SDS-PAGE. Ni-NTA chromatography.


Endotoxin: <0.1 EU per 1 µg of the protein by the LAL method.

Amino acid sequence:

VLETNDTNLKCQCLRSTSNWVPIRLIEKIQIWPPGNGCPTREVIVWMTNKAICLN PQSKLLQKLINLMWRKKTSTTL PAVSKKSIA with

polyhistidine tag at the N-terminus.

Formulation: Lyophilized from a sterile filtered aqueous solution in 1×PBS, pH 7.4.



SDS-PAGE analysis of recombinant swine CXCL13

Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H₂O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.