

Recombinant Mouse, IL36 γ / Interleukin-36 gamma,

Animal Free & Carrier Free

Cat #: C-CYP190

Size: 5 μ g, 20 μ g, 100 μ g, 500 μ g, 1mg

Shipping: Blue Ice

Product Overview

The expression of IL-36 γ is high in psoriatic plaques and in tissues including epithelial cells. Signaling of IL-36 γ through the IL-1Rrp2 receptor, which is mainly expressed on certain dendritic cells. The interaction between the IL-36 ligands and IL-1Rrp2 receptor trigger dendritic cell maturation and activation. IL-36 γ also work as an agonist of NF- κ B, consequently, stimulate the inflammatory response in bronchial epithelial cells.

Product Information

Source: *Escherichia Coli*.

Purity: >98% as determined by SDS-PAGE. Ni-NTA chromatography.

Biological Activity: Measure by its ability to induce IL-6 secretion in 3T3 cells. The ED₅₀ for this effect is < 15 ng/mL.

The specific activity of recombinant mouse IL-36 gamma is > 6 x 10⁴ IU /mg.


Endotoxin: <0.1 EU per 1 μ g of the protein by the LAL method.

Amino acid sequence:

MGRETPDFGEVFDLDQQVWIFRNQALVTVPRSHRVTPVSVTILPCKYPESLEQDKGIAIYLGIQNPDKCLFCKEVNGHPTLLLKEEKILDLYH

HPEPMKPFLFYHTRTGGTSTFESVAFPGHYIASSKTGNPIFLTSTKKGEEYNNINFNLDIKS with polyhistidine tag at the C-terminus.

Formulation: Lyophilized from a sterile filtered aqueous solution in 1 \times PBS, pH 7.4.



SDS-PAGE analysis of recombinant mouse IL-36 gamma

Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H₂O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.