

# Recombinant Mouse, IL38 / Interleukin-38, Animal Free & Carrier Free

Cat #: C-CYP196

Size: 5μg, 20μg, 100μg, 500μg, 1mg

Shipping: Blue Ice

### **Product Overview**

IL-38 expresses in the fetal skin, spleen and tonsil, mostly in the basal epithelia of skin and in proliferating B cells of the tonsil. IL-38 can bind soluble IL-1 receptor type 1 and may be involved in the regulation of adapted and innate immune responses.

## **Product Information**

Source: Escherichia Coli.

**Purity:** >98% as determined by SDS-PAGE. Ni-NTA chromatography.

**Endotoxin:** <0.1 EU per 1  $\mu g$  of the protein by the LAL method.

Amino acid sequence:


MCSLPMARYYIIKDAHQKALYTRNGQLLLGDPDSDNYSPEKVCILPNRGLDRSKVPIFLGMQGGSCCLACVKTREGP

LLQLEDVNIEDLYKGGEQTTRFTFFQRSLGSAFRLEAAACPGWFLCGPAEPQQPVQLTKESEPSTHTEFYFEMSR with polyhistidine tag at the C-terminus.

Formulation: Lyophilized from a sterile filtered aqueous solution in 1×PBS, pH 7.4.







SDS-PAGE analysis of recombinant mouse IL-38

## **Usage Method**

- 1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
- 2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled  $H_2O$ . Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
- 3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
- 4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10  $\mu$ g/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

## **Storage**

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

#### **Note**

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.

