

# Recombinant Mouse/Rat TGFβ1 / Transforming Growth Factor beta 1, Animal Free & Carrier Free

Cat #: C-CYP235

Size: 5µg, 20µg, 100µg, 500µg, 1mg

Shipping: Blue Ice

## Product Overview

Transforming growth factor beta 1 or TGF-β1 is a polypeptide member of the transforming growth factor beta superfamily of cytokines. It is a secreted protein that performs many cellular functions, including the control of cell growth, cell proliferation, cell differentiation, and apoptosis. In humans, TGF-β1 is encoded by the TGFB1 gene.

## Product Information

**Source:** *Escherichia Coli*.

**Purity:** >98% as determined by SDS-PAGE. Ni-NTA chromatography.


**Biological Activity:** Measure by its ability to inhibit the IL-4 dependent proliferation in HT-2 cells. The ED<sub>50</sub> for this effect is < 0.1 ng/mL.

**Endotoxin:** <0.1EU per 1µg of the protein by the LAL method.

**Amino acid sequence:**

MALDTNYCFSSTEKNCCVRQLYIDFRKDLGWKWIHEPKGYHANFCLGPCPYIWSLDTQYSKVLALYNQHNP GASASPCCV PQALEPLIVY YVGRKPKVEQLSNMIVRSC KCS with polyhistidine tag at the C-terminus.

**Formulation:** Lyophilized from a sterile filtered aqueous solution in 20 mM sodium citrate, 0.2 M NaCl, pH 3.5.



SDS-PAGE analysis of recombinant mouse TGF beta 1

## Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H<sub>2</sub>O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

## Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

## Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.