

Recombinant Human TRAIL / TNF-Related Apoptosis-Inducing Ligand, Animal Free & Carrier Free

Cat #: C-CYP240

Size: 5µg, 20µg, 100µg, 500µg, 1mg

Shipping: Blue Ice

Product Overview

TRAIL acts as a cytotoxic protein, through activating rapid apoptosis in tumor cells (not in normal cells). TRAIL conducts apoptosis through binding to DR4 and DR5, which are two death-signaling receptors belonging to the TNFR superfamily of transmembrane proteins. These receptor contain a cytoplasmic " death domain", which can initiate the cell's apoptotic process.

Product Information

Source: *Escherichia Coli*.

Purity: >98% as determined by SDS-PAGE. Ni-NTA chromatography.


Biological Activity: Measure by its ability to induce cytotoxicity in L929 cells in the presence of actinomycin D. The ED₅₀ for this effect is 10.4-15.4 ng/mL.

Endotoxin: <0.1EU per 1µg of the protein by the LAL method.

Amino acid sequence:

MRERGPQRVAAHITGTRGRSNTLSSPNSKNEKALGRKINSWESSRSGHSFLSNLHLRNGELVIHEKGFYYIYSQTYFRFQEEIKENTKNDKQ
MVQYIYKYTSYPDPILLMKSARNSCWSKDAEYGLYSIQGGIFELKENDRIFVSVTNEHLIDMDHEASFFGAFLVG with polyhistidine tag at the C-terminus.

Formulation: Lyophilized from a sterile filtered aqueous solution in 1X PBS, pH 8.0.



SDS-PAGE analysis of recombinant human TRAIL

Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H₂O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

Storage

| Physical Appearance | Storage | Stability |
|---------------------------------|----------------|------------------|
| Lyophilized powder | -20°C to -80°C | 1 year |
| Reconstitution (initial) | 2°C to 8°C | Less than 1 week |
| Reconstitution (after dilution) | -20°C to -80°C | 3 to 6 months |

Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.