

## Recombinant Mouse CCL3 / C-C Motif Chemokine Ligand 3,

### Animal Free & Carrier Free

Cat #: C-CYP266

Size: 5µg, 20µg, 100µg, 500µg, 1mg

Shipping: Blue Ice

### Product Overview

CCL3 is belonging to the CC chemokine family. CCL3 participates in activating and recruiting cells, such as lymphocytes, monocytes, and granulocytes during acute inflammation. In addition, CCL3 can enhance IFN- $\gamma$  secretion from activated T cells and thus induces Th1 response, thereby to regulating leukocyte migration. It is reported that CCL3 is involved in susceptibility of HIV infection and disease progression of AIDS.

### Product Information

**Source:** *Escherichia Coli*.

**Purity:** >98% as determined by SDS-PAGE. Ni-NTA chromatography.


**Biological Activity:** Measure by its ability to chemoattract human PBMCs using a concentration range of 10 - 100 ng/mL.

Note: Results may vary from different PBMC donors.

**Endotoxin:** <0.1EU per 1µg of the protein by the LAL method.

**Amino acid sequence:** APYGADTPTACCFYSYRKIPRQFIVDYFETSSLCSQPGVIFLTKRNRQICADSKETWVQEYITDLELNA with polyhistidine tag at the N-terminus.

**Formulation:** Lyophilized from a sterile filtered aqueous solution in 1X PBS, pH 7.4



SDS-PAGE analysis of recombinant mouse CCL3

## Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H<sub>2</sub>O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

## Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

## Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.