

## Recombinant Human BMP12 / Bone Morphogenetic Protein 12, Animal Free & Carrier Free

Cat #: C-CYP267

Size: 5µg, 20µg, 100µg, 500µg, 1mg

Shipping: Blue Ice

### Product Overview

Bone morphogenetic proteins (BMPs) are a group of growth factors also known as cytokines and as metabologens.

BMP-12 regulates chondrogenesis, bone morphogenesis, and neuron differentiation.

### Product Information

**Source:** *Escherichia Coli*.

**Purity:** >98% as determined by SDS-PAGE. Ni-NTA chromatography.

**Biological Activity:** Measure by its ability to induce alkaline phosphatase production by ATDC5 cells. The ED<sub>50</sub> for this effect is < 2 µg/mL.


**Endotoxin:** <0.1EU per 1µg of the protein by the LAL method.

**Amino acid sequence:**

MTALAGTRTAQGGGGAGRGHGRRGRSRKPLHVDFKELGWDDWIIAPLDYEAYHCEGLCDFPLRSHLEPTNHAIQTLNMAPDA

APASCCVPARLSPISILYIDAANNVVYKQYEDMVVEACGCR with polyhistidine tag at the C-terminus.

**Formulation:** Lyophilized from a sterile filtered aqueous solution in 20 mM sodium citrate, 0.2 M NaCl, pH 3.5



SDS-PAGE analysis of recombinant human BMP12

## Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H<sub>2</sub>O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

## Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

## Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.