

Recombinant Human FGF9 / Fibroblast Growth Factor 9,

Animal Free & Carrier Free

Cat #: C-CYP312

Size: 5µg, 20µg, 100µg, 500µg, 1mg

Shipping: Blue Ice

Product Overview

FGF9 (fibroblast growth factor-9), also called HBGF-9 (heparin-binding growth factor-9) and GAF (glia-activating factor), is an approximately 26 kDa secreted glycoprotein of the FGF family (1-3). FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. FGF9 plays an important role in the regulation of embryonic development, cell proliferation, cell differentiation and cell migration.

Product Information

Source: *Escherichia Coli*.

Purity: >95% as determined by SDS-PAGE. Ni-NTA chromatography.


Biological Activity: Measure by its ability to induce 3T3 cells proliferation. The ED₅₀ for this effect is <2 ng/mL.

Endotoxin: <0.1 EU per 1 µg of the protein by the LAL method.

Amino acid sequence:

MPLGEVGNVYFGVQDAVPFGNVPVLPVDSPVLLSDHLGQSEAGGLPRGPAVTDLDHLKILRRRQLYCRTGFHLEIFPNGTIQGTRKDHSRF
GILEFISIAVGLVSIRGVDSGLYLGMNEKGELYGSEKLTQECVFREQFEENWYNTYSSNLYKHVDTGRRYYVALNKDGTREGTRTRKRHQKFT
H FLPRVDPDKVPELYK DILSQS with polyhistidine tag at the C-terminus.

Formulation: Lyophilized from a sterile filtered aqueous solution in 1×PBS, pH 7.4.



SDS-PAGE analysis of recombinant human FGF9

Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H₂O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.