

Recombinant Human FGF12 / Fibroblast Growth Factor 12,

Animal Free & Carrier Free

Cat #: C-CYP317

Size: 5µg, 20µg, 100µg, 500µg, 1mg

Shipping: Blue Ice

Product Overview

Fibroblast growth factor 12 (FGF12) is a member of the FGF superfamily of molecules which contains at least 22 members. FGF family members possess broad mitogenic and cell survival activities and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth, and invasion. The specific function of FGF12 gene has not yet been determined. Two alternatively spliced transcript variants encoding distinct isoforms have been reported.

Product Information

Source: *Escherichia Coli*.

Purity: >98% as determined by SDS-PAGE. Ni-NTA chromatography.

Biological Activity: Measure by its ability to induce 3T3 cells proliferation. The ED₅₀ for this effect is <2 ng/mL.

Endotoxin: <0.1 EU per 1 µg of the protein by the LAL method.


Amino acid sequence:

MESKEPQLKGIVTRLFSQQGYFLQMHPDGTIDGTDKENS DYTLFNLIPVGLRVVAIQGVKASLYVAMNGEGYLYSSDVFTPECKFKESVFEN

YYVIYSSTLYRQQESGRAWFLGLNKEGQIMKGNRVKKT KPSSHFPKPIEVCMYREPSLHEIGEKQGRSRKSSGTPTMNGGKVVNQDST

with polyhistidine tag at the C-terminus.

Formulation: Lyophilized from a sterile filtered aqueous solution in 1×PBS, pH 7.4.



SDS-PAGE analysis of recombinant human FGF12

Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H₂O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.