

# Recombinant Human CDNF / Cerebral Dopamine Neurotrophic Factor, Animal Free & Carrier Free

Cat #: C-CYP351

Size: 5μg, 20μg, 100μg, 500μg, 1mg

Shipping: Blue Ice

#### **Product Overview**

Cerebral dopamine neurotrophic factor also known as ARMET-like protein 1 or is a protein that in humans that is encoded by the CDNF gene. CDNF protein is expressed in human brain, acts differently from known neurotrophic factors and can protect and repair dopamine neurons in two pre-clinical models of Parkinson's disease (PD).

#### **Product Information**

Source: Escherichia Coli.

Purity: >98% as determined by SDS-PAGE. Ni-NTA chromatography.

**Endotoxin:** <0.1EU per 1µg of the protein by the LAL method.

#### Amino acid sequence:

MQEAGGRPGADCEVCKEFLNRFYKSLIDRGVNFSLDTIEKELISFCLDTKGKENRLCYYLGATKDAATKILSEVTRPMSVHMPAMKICEKLKK LDSQICELKYEKTLDLASVDLRKMRVAELKQILHSWGEECRACAEKTDYVNLIQELAPKYAATHPKTEL with polyhistidine tag at the C-terminus.

Formulation: Lyophilized from a sterile filtered aqueous solution in 1×PBS, pH 8.0.







SDS-PAGE analysis of recombinant human CDNF

## **Usage Method**

- 1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
- 2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled  $H_2O$ . Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
- 3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
- 4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10  $\mu$ g/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

### **Storage**

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

#### **Note**

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.

