

Recombinant Human, Gal-13 / Galectin-13,

Animal Free & Carrier Free

Cat #: C-CYP372

Size: 5µg, 20µg, 100µg, 500µg, 1mg

Shipping: Blue Ice

Product Overview

Galectins are a class of proteins that bind specifically to β -galactoside sugars. There have been 15 galectins discovered in mammals, encoded by the LGALS genes. Only galectin-1, -2, -3, -4, -7, -8, -9, -10 and -12 have been identified in humans. Galectin-13, also known as placental protein 13, is a placenta-specific galectin that induces the apoptosis of T lymphocytes, which may reduce the danger of maternal immune attacks on the fetal semiallograft during the long gestation of anthropoid primates.

Product Information

Source: *Escherichia Coli*.

Purity: >95% as determined by SDS-PAGE. Ni-NTA chromatography.


Endotoxin: <0.1EU per 1µg of the protein by the LAL method.

Amino acid sequence:

SSLPVPYKLPVSLSVGSCVIIKGTPIHSFINDPQLQVDFYTDMEEDSDIAFRFRVHFGNHVVMNRRREFGIWMLEETTDYVPFEDGKQFELCI

YVHYNEYEIKVNGIRIYGFVHRIPPSFVKMVQVSRDISLTSVCVCN with polyhistidine tag at the N-terminus.

Formulation: Lyophilized from a sterile filtered aqueous solution in 1xPBS, pH 7.4.



SDS-PAGE analysis of recombinant human Galectin-13

Usage Method

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H₂O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 µg/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.