

# Recombinant Human, Gal-14 / Galectin-14,

# **Animal Free & Carrier Free**

Cat #: C-CYP373 Size: 5µg, 20µg, 100µg, 500µg, 1mg Shipping: Blue Ice

### **Product Overview**

Galectin14 is a member of the galectin family, of which members all contain one or two carbohydrate recognition domains targeting the beta-galactoside. The LGALS14 gene is predominantly expressed in the placenta. The gene encoded protein is identified inside the cells of placenta and eosinophils. LGALS14 is released by eosinophils following allergen stimulation.

### **Product Information**

Source: Escherichia Coli.

Purity: >98% as determined by SDS-PAGE. Ni-NTA chromatography.

**Endotoxin:** <0.1EU per  $1\mu g$  of the protein by the LAL method.

Amino acid sequence:

SSLPVPYTLPVSLPVGSCVIITGTPILTFVKDPQLEVNFYTGMDEDSDIAFQFRLHFGHPAIMNSCVFGIWRYEEKCYYLPFEDGKPFELCIYVR HKEYKVMVNGQRIYNFAHRFPPASVKMLQVFRDISLTRVLISD with polyhistidine tag at the N-terminus.

Formulation: Lyophilized from a sterile filtered aqueous solution in 1×PBS, pH 7.4.





kDa	М
75 63 48	
35	
25	Same P
17 10	-

SDS-PAGE analysis of recombinant human Galectin-14

### **Usage Method**

1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.

2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H<sub>2</sub>O. Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.

3. The reconstituted solution can be stored at 2-8°C for up to 1 week.

4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10  $\mu$ g/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

#### Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

#### Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.

