

Recombinant Human, CXCL1 / C-X-C Motif Chemokine Ligand 1, Animal Free & Carrier Free

Cat #: C-CYP376

Size: 5μg, 20μg, 100μg, 500μg, 1mg

Shipping: Blue Ice

Product Overview

CXCL1/GROa act as a growth factor for melanoma cells. It is also a chemotaxin for neutrophils. Like other alpha chemokines, this protein is potent neutrophil attractant and activator and is also affect basophils. Clinical research show that CXCL1 protein may be a therapeutic target as well as a diagnostic marker in ovarian cancer.

Product Information

Source: Escherichia Coli.

Purity: >98% as determined by SDS-PAGE. Ni-NTA chromatography.

Biological Activity: chemoattract BaF3 cells transfected with human CXCR2. The ED50 for this effect is <3 ng/mL.

Endotoxin: <0.1EU per $1\mu g$ of the protein by the LAL method.


Amino acid sequence: NVKSPGPHCAQTEVIATLKNGRKACLNPASPIVKKIIEKMLNSDKSN with polyhistidine tag at the

N-terminus.

Formulation: Lyophilized from a sterile filtered aqueous solution in 1×PBS, pH 7.4.







SDS-PAGE analysis of recombinant human CXCL1

Usage Method

- 1. Before opening, it is recommended to centrifuge at 3000-3500 rpm for 5 minutes.
- 2. Reconstitute to a concentration of 0.1-1.0 mg/mL in sterile distilled H_2O . Allow the solution to sit at room temperature for at least 20 minutes to ensure complete dissolution. Avoid vigorous vortexing.
- 3. The reconstituted solution can be stored at 2-8°C for up to 1 week.
- 4. For long-term storage, it is recommended to further dilute the solution with a carrier protein (such as 0.1% BSA, 10% FBS, or 5% HSA) to a concentration of no less than 10 μ g/mL and aliquot for storage at -20°C to -80°C for 3 to 6 months. If serum-free experiments are required, a 5% trehalose solution can be used as a carrier instead. Avoid repeated freeze-thaw cycles.

Storage

Physical Appearance	Storage	Stability
Lyophilized powder	-20°C to -80°C	1 year
Reconstitution (initial)	2°C to 8°C	Less than 1 week
Reconstitution (after dilution)	-20°C to -80°C	3 to 6 months

Note

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.

